

## **Branched and linear forms of PFAS – A means of a more comprehensive assessment of environmental impacts**

*Patrick van Hees, Eurofins Environment Testing Sweden AB*

### **Abbreviations:**

ECF = Electrochemical fluorination; FTOH = Fluorotelomer alcohols; N-EtFOSA = N-ethylperfluorooctanesulfonamide; N-EtFOSE = N-ethylperfluorooctanesulfonamidoethanol; N-MeFOSA = N-methylperfluorooctanesulfonamide; N-MeFOSE = N-methylperfluorooctanesulfonamidoethanol; PFAS = Poly- and perfluoroalkyl substances; PFCA = perfluorinated carboxylic acids; PFHxS = Perfluorohexane sulfonate; PFOA = Perfluorooctanoate; PFOS = Perfluorooctane sulfonate; PFOSA = Perfluorooctanesulfonamide

### *Introduction*

Over the last 20-25 years the research on both environmental and health related effects of PFAS has been intense. During the last 5-10 years the difference between isomeric forms of the substances has emerged as an additional factor. The physicochemical properties of the linear and branched forms vary slightly which leads to significant differences in biochemical reactions/transformations, bioaccumulation and thus potential toxic exposure. Adsorption onto solid phases is another property affected. All these factors also influence the relative distribution in environmental compartments such as water, soil and sediment. The research has also created an opportunity to develop environmental forensic applications.

### *Manufacturing processes*

Electrochemical fluorination (ECF) and telomerisation are the two major methods employed to produce PFAS. The branched isomers of PFAS are mainly manufactured in the ECF method, which has historically been used to produce the major part of the two dominant PFAS, PFOS and PFOA. ECF gives rise to complex mixtures of linear and branched compounds. PFOA produced by this method has typically had an isomer composition of 78% linear (n-PFOA) and 22% branched isomers (br-PFOA). ECF-PFOS shows a distribution of around 70% linear (n-PFOS) and 30% branched (br-PFOS). Precursors and similar substances e.g. N-methyl and N-ethyl perfluorooctane sulfonamido ethanols (Me- and EtFOSEs) and sulfonamides (Me- and EtFOSAs) have ratios in the same range. On the other hand the telomerisation process keeps the structure of the starting telogen and a pure linear or isopropyl form is produced (Benskin et al., 2010; Jiang et al, 2015).

### *Distribution in environmental matrices*

The differing properties of linear and branched PFAS isomers can affect the partitioning and migration of the substances in the aquatic environment. A number of studies have reported enrichment of br-PFOS in natural waters as compared to technical products (Benskin et al., 2010; Chen et al., 2015 and references within).

In a nationwide study in France it was found that n-PFOS was the dominant isomer in sediments (88% of total; n=129), while in water n-PFOS comprised 48% (n=333) (Munoz et al., 2015). A similar distribution was observed by Houde et al. (2008). Chen et al. (2015) studied PFOS, PFOA and PFOSA in water, water particulate phase and sediment. For PFOS and PFOSA there was more of the linear form adsorbed to the solid phases than found in water (PFOS 70-85% vs 40-43%; PFOSA: 93-95% vs 64-75%). For PFOA the difference was not as large in absolute numbers but calculations still indicated a higher log  $K_{OC}$  for n-PFOA.

In a study by Kärman et al. (2011) soil from a fire-fighting training platform contained 63% n-PFOS but this fraction had increased to 80-85% 200m away from the site. In the receiving lake the sediment was also enriched in the linear form (75-85%). However, in the seepage water from the site the opposite was observed where n-PFOS accounted for a little less than 60%. PFOA showed a similar pattern. At the source 80% was in the form of n-PFOA in the soil, which rose to 100% within 75m distance. In seepage the corresponding figure was 75%. PFOA was not found in the sediment.

In an investigation of three Swedish sites drinking water, untreated and treated waste water and sewage sludge, branched and linear forms of PFOS, PFOSA, PFHxS and PFOA were determined (Filipovic and Berger, 2015). The contribution of n-PFOS was on average 62% in the untreated waste water while in the sludge was 86%. Also for PFOSA a tendency for a higher percentage of the linear form in the sludge as compared to influent water was observed (83% vs 68%). For PFOA >91% was linear in the untreated water and in the sludge no branched forms could be detected. Results for PFHxS were not conclusive since all sludge samples had concentrations below limit of detection.

Also for water cleaning the isomeric profile can be important. Östlund (2015) investigated PFAS removal in pilot scale column experiments using anion exchange resin and granular activated carbon. The adsorption efficiency of branched and linear PFOS, PFOSA and PFHxS was assessed. For the anion exchanger no difference due to isomeric form was found, while for active carbon a lower removal of the branched form was seen. At the end of the study (33000 bed volumes) the difference in adsorption was 86% vs 78% for PFOS. The same numbers were 69% vs 57% for PFOSA and 57% vs 40% for PFHxS.

### *Source tracking*

Comparison of the isomeric forms of PFAS in environmental samples with commercial products may yield insights to the manufacturing origins (Benskin et al., 2010). Chen et al. (2015) used an isomer profiling technique to calculate the relative contributions of various industrial origins for PFOA in a Chinese river. The outcome indicated that the major source was electrochemical fluorination (ECF) (55%), followed by linear telomerisation (41%) and isopropyl telomerisation (4%).

Another field of research is to what extent the isomeric composition in e.g. water reflects the transformation of precursors, and the possibility that branched forms are more prone in this regard (Benskin et al, 2010). Leaching tests of sediments showed that the fraction of n-PFOS in the leachate was much higher than in the water in the field (Chen et al., 2015). The

difference between the leaching and field results suggested that PFOS precursors could make a contribution to the linear/branched ratio of PFOS in the natural system. A similar observation was made by Kärrman et al (2011) where PFOS in soil in the source area contained less linear PFOS (63%) than could be expected from ECF products.

Benskin et al (2011) used the absence of branched perfluorinated carboxylic acids (PFCA) as one line of evidence that the compounds observed (PFOA and longer homologues) originated from oxidation of above all FTOH rather direct transport (vapour phase and particles) of PFCAs.

#### *Isomer distribution in humans and biota*

For PFAS four major pathways for direct and indirect exposure can be outlined: ingestion of dust, dietary and drinking water intake, and inhalation of air (Gebbinck et al., 2015). Concerning food, n-PFOS comprised 92% of total PFOS in a general Swedish diet and estimates for PFOSA indicated about 98% of the linear form. PFOS and PFOSA isomer ratios in drinking water were 60% n-PFOS and 58% n-PFOSA. Considering total PFOS exposure, taking all pathways into account, the proportion was estimated as 84% linear and 16% branched. The number could be dependent upon the diet, especially at high fish consumption (Gebbinck et al., 2015). Enrichment of the linear form of PFOS in animal tissue is frequently seen (often >90% n-PFOS) e.g. for fish and pelagic organisms as well as mammals such as rats and polar bears (Kärrman et al., 2011; Greaves and Letcher, 2013; Miralles-Marco and Harrad, 2015)

However, in humans the isomeric PFOS profile may be more complicated recent studies suggest (Miralles-Marco and Harrad, 2015). In particular human serum samples can sometimes show a br-PFOS content even higher than in the technical ECF products. It has frequently been put forward that these observations can be explained by either an inaccurate estimation of exposure (e.g., unknown precursors, missing or poorly quantified pathways and/or isomer profiles of PFOS and precursors) or an incomplete understanding of the human pharmacokinetic processes (e.g. biotransformation and elimination kinetics) (Gebbinck et al., 2015).

Also exposure to PFAS contamination can alter the isomeric profile in humans. In a study of young women in Uppsala (SE) higher total levels of PFHxS as well as an increased fraction of br-PFHxS in serum were observed for those who had been exposed to contaminated drinking water. Still the percentage of br-PFHxS in serum was lower than that found in water (6 vs 20%) (Gyllenhammar et al., 2015).

#### *References*

- Benskin, J.P., De Silva, A. O., Martin, J. W. Isomer Profiling of Perfluorinated Substances as a Tool for Source Tracking: A Review of Early Findings and Future Applications. *Rev. Environ Contam. Toxicol.*, 2010, 208, 111-160
- Benskin, J.P., Phillips, V., St. Louis V.L., Martin, J.W. Source Elucidation of Perfluorinated Carboxylic Acids in Remote Alpine Lake Sediment Cores, *Env. Sci. Tech.*, 2011, 45, 7188–7194
- Chen X., Zhu, L., Pan, X., Fang, S., Zhang, Y., Yang, L. Isomeric specific partitioning behaviors of perfluoroalkyl substances in water dissolved phase, suspended particulate matters and sediments in Liao River Basin and Taihu Lake, China. *Wat Res.* 2015, 80, 235-244

- Filipovic, M., Berger, U. Are perfluoroalkyl acids in waste water treatment plant effluents the result of primary emissions from the technosphere or of environmental recirculation? *Chemosphere* 2015, 129, 74–80
- Gebbink, W. A., Berger, U., Cousins, I. T. Estimating human exposure to PFOS isomers and PFCA homologues: The relative importance of direct and indirect (precursor) exposure. *Env. Int.* 2015, 74, 160–169
- Greaves, A. K., Letcher, R.J. Linear and branched perfluorooctane sulfonate (PFOS) isomer patterns differ among several tissues and blood of polar bears. *Chemosphere* 2013, 93, 574–580
- Gyllenhammar I., Berger, U., Sundström, M., McCleaf, P., Eurén, K., Eriksson, S., Ahlgren, S., Lignell, S., Aune, M., Kotova, N., Glynn, A. Influence of contaminated drinking water on perfluoro alkyl acid levels in human serum – A case study from Uppsala, Sweden. *Env. Res.* 2015, 140, 673–683.
- Houde, M., Jecub, M., Small, J., Backus, S., Owawang, X., Alae, L., Derekc, A., Muir, U. Fractionation and Bioaccumulation of Perfluorooctane Sulfonate (PFOS) Isomers in a Lake Ontario Food. *Environ. Sci. Technol.* 2008, 42, 9397–9403
- Jiang, Y., Zhang, L., Yang, X, Chu, L., Zhu, W. Perfluoroalkyl acids (PFAAs) with isomer analysis in the commercial PFOS and PFOA products in China. *Chemosphere* 2015, 127, 180–187
- Kärman, A., Elgh-Dalgren, K., Lafossas A. Mørskeland T., Environmental levels and distribution of structural isomers of perfluoroalkyl acids after aqueous fire-fighting foam (AFFF) contamination. *Environ. Chem.* 2011, 8, 372–380
- Miralles-Marco, A., Harrad, S. Perfluorooctane sulfonate: A review of human exposure, biomonitoring and the environmental forensics utility of its chirality and isomer distribution. *Env. Int.* 2015, 77, 148–159
- Munoz, G., Giraudel, J-L., Botta, F., Lestremau, F., Dévier, M-H., Budzinski, H., Labadie, P. Spatial distribution and partitioning behavior of selected poly- and perfluoroalkyl substances in freshwater ecosystems: A French nationwide survey. *Sci. Tot. Environ.* 2015, 517, 48–56
- Östlund, A. Removal Efficiency of Perfluoroalkyl Substances (PFASs) in Drinking Water – Evaluation of granular activated carbon (GAC) and anion exchange (AE) using column tests, and the effect of dissolved organic carbon. MSc thesis, Swe Univ Agricult. Sci., Uppsala, 2015